

	Week 1	Week 2	Week 3 (maintenance)
Weight (kg)	Volume administered twice a day	Volume administered twice a day	Volume administered twice a day
2 to <2.5	0.3 mL	0.6 mL	0.8 mL
2.5 to <3	0.4 mL	0.8 mL	1 mL
3 to <3.5	0.5 mL	0.9 mL	1.2 mL
3.5 to <4	0.5 mL	1.1 mL	1.4 mL
4 to <4.5	0.6 mL	1.2 mL	1.6 mL
4.5 to <5	0.7 mL	1.4 mL	1.8 mL
5 to <5.5	0.8 mL	1.5 mL	2 mL
5.5 to <6	0.8 mL	1.7 mL	2.2 mL
6 to <6.5	0.9 mL	1.8 mL	2.4 mL
6.5 to <7	1.0 mL	2.0 mL	2.6 mL
7 to <7.5	1.1 mL	2.1 mL	2.8 mL
7.5 to <8	1.1 mL	2.3 mL	3 mL
8 to <8.5	1.2 mL	2.4 mL	3.2 mL
8.5 to <9	1.3 mL	2.6 mL	3.4 mL
9 to <9.5	1.4 mL	2.7 mL	3.6 mL
9.5 to <10	1.4 mL	2.9 mL	3.8 mL
10 to <10.5	1.5 mL	3.0 mL	4 mL
10.5 to <11	1.6 mL	3.2 mL	4.2 mL
11 to <11.5	1.7 mL	3.3 mL	4.4 mL
11.5 to <12	1.7 mL	3.5 mL	4.6 mL
12 to <12.5	1.8 mL	3.6 mL	4.8 mL

- **Administer twice daily** at least 9 hours apart
- **Reduce the risk of hypoglycemia by administering Hemangeol** during or right after a feeding. **Skip the dose** if child is not eating or is vomiting.
- **Monitor heart rate and blood pressure for 2 hours** after Hemangeol initiation or dose increases.