

INSTRUCTIONS FOR USE
HEMANGEOL® (he-man je-ohl)
(propranolol hydrochloride oral solution)


Read these Instructions for Use before giving a dose of HEMANGEOL to your child for the first time and each time you get a refill. There may be new information. Your doctor or pharmacist should show you how to correctly measure and give a dose of HEMANGEOL to your child before you give it for the first time.

Important: Read the Medication Guide that comes with HEMANGEOL.

- To reduce the risk of your child getting low blood sugar (hypoglycemia), you must give HEMANGEOL either during a feeding or right away after a feeding.
- **Do not give a dose of HEMANGEOL if your child is vomiting, is not taking feedings, or is showing signs or symptoms of low blood sugar.**

When you get HEMANGEOL from your doctor or pharmacist, you will receive a box that contains the supplies needed to give HEMANGEOL to your child, including:

- One glass bottle of HEMANGEOL
- One 5 mL oral dosing syringe (inside a plastic bag) that is marked to help you correctly measure a dose of HEMANGEOL (See Figure A). If the carton does not contain the oral dosing syringe, ask your pharmacist to give you an oral dosing syringe that can be used to measure HEMANGEOL.




Preparing to give your child a dose of HEMANGEOL:

Step 1. Place your box of supplies on a clean flat work surface, such as a table.


Step 2. Remove the HEMANGEOL bottle and oral dosing syringe from the box (See Figure A above). **Do not shake the bottle before use.** Keep the box for storage.

Step 3. Remove the oral dosing syringe from the plastic bag. Safely throw the plastic bag away. The barrel of the syringe has markings in milliliters (mL). Look at the markings on the barrel of the oral dosing syringe and find the mL marking that matches the HEMANGEOL dose in mL prescribed by your doctor (See Figure B).




Step 4. Open the bottle of HEMANGEOL by pushing down on the plastic cap while turning the cap to the left (See Figure C).

- Write down on the box the date when you first open the bottle.




Step 5. Place the bottle on your work surface. Use one hand to hold the bottle upright. Use your other hand to insert the tip of the oral dosing syringe into the syringe adapter at the top of the bottle. Push the plunger all the way down (See Figure D).


- Do not remove the syringe adapter. If the syringe adapter is missing talk to your pharmacist.




Step 6. Use one hand to hold the oral dosing syringe in place. With your other hand, turn the bottle upside down. Pull back on the plunger until the top of the plunger lines up with the marking on the barrel of the syringe that matches the dose of HEMANGEOL prescribed by your doctor (See Figure E). Your child's dose may be different than the dose shown in Figure E.



Step 7. Check for air bubbles in the oral dosing syringe. If you see air bubbles, push up on the plunger towards the bottle just enough to remove any large air bubbles and then pull back to the measured dose (See Figure F).




Step 8. Turn bottle upright again and place it in on your work surface. Remove the oral dosing syringe from the bottle (See Figure G). Do not push the plunger in during this step. The syringe adapter should stay attached to the bottle.




Giving your child a dose of HEMANGEOL:

Step 9. Slowly squirt HEMANGEOL into your child's mouth after placing the oral dosing syringe against the inside of the cheek (See Figure H).

- Keep your child in an upright position for a few minutes right after giving a dose of HEMANGEOL.
- If needed, you can dilute the dose of HEMANGEOL in a small amount of milk or fruit juice and give it to your child in a baby's bottle. If your child spits up a dose or if you are not sure your child got all of the medicine, do not give another dose. Wait until the next scheduled dose.




Step 10. Replace the plastic cap on the bottle. Close the bottle by turning the plastic cap to the right (See Figure I).



Cleaning the oral dosing syringe:

Step 11. Clean the oral dosing syringe after each use by rinsing with clean tap water (See Figure J).

- Do not take apart the oral dosing syringe.
- Do not use any soap or alcohol based product to clean. Wipe the outside dry.
- Do not put the oral dosing syringe through a sterilizer or dishwasher.



Step 12: Place the bottle and the oral dosing syringe in the box.

How should I store HEMANGEOL?

- When not in use, keep the bottle of HEMANGEOL and the oral dosing syringe in the box it comes in.
 - Store HEMANGEOL at room temperature, between 68 °F to 77 °F (20 °C to 25 °C). Do not freeze.
 - Safely throw away any opened bottle of HEMANGEOL after 2 months, even if there is medicine left in the bottle.
- Keep HEMANGEOL and all medicines out of the reach of children.**

This Instructions for Use has been approved by the U.S. Food and Drug Administration.